

On the Nörlund-Rice Integral Formula

R. Sivaraman, M. Muniru Iddrisu, J. López-Bonilla



Abstract: After introducing the famous Nörlund-Rice integral formula, we apply it to Laguerre polynomials, Melzak's relation, and Stirling numbers of the second kind to obtain nice expressions.

Keywords: Stirling numbers, Nörlund-Rice integral, Melzak's identity, Laguerre polynomials.

I. INTRODUCTION

We have the Nörlund-Rice integral formula [1-6]:

$$\sum_{k=0}^n \binom{n}{k} (-1)^k f(k) = \frac{(-1)^n n!}{2\pi i} \oint_C \frac{f(z) dz}{z(z-1)(z-2)\cdots(z-n)}, \quad n \geq 0, \quad (1)$$

where f is analytic in a domain containing the interval $[0, n]$, and C is a closed, simple, positively oriented curve surrounding $[0, n]$.

In Sec. 2, we provide the applications of (1) to Stirling numbers of the second kind, Melzak's identity, and Laguerre polynomials.

II. APPLICATIONS OF NÖRLUND-RICE'S FORMULA

a).- Let us consider the analytic function $f(z) = z^m$, $m \geq 0$ and apply it to (1).

Then (1) gives interesting expressions for the Stirling numbers of the second kind [7-9][20][21][22]:

$$S_m^{[n]} = \frac{(-1)^n}{n!} \sum_{k=0}^n \binom{n}{k} (-1)^k k^m = \frac{1}{2\pi i} \oint_C \frac{z^m dz}{z(z-1)\cdots(z-n)}, \quad m, n \geq 0, \quad (2)$$

thus:

$$S_0^{[0]} = \frac{1}{2\pi i} \oint_C \frac{dz}{z} = 1, \quad S_m^{[0]} = \frac{1}{2\pi i} \oint_C z^{m-1} dz = 0, \quad m \geq 1, \quad S_m^{[1]} = \frac{1}{2\pi i} \oint_C \frac{z^{m-1}}{z-1} dz = \begin{cases} 0, & m = 0, \\ 1, & m \geq 1, \end{cases}$$

A. Theorem

The Stirling's numbers of the second kind satisfy the recurrence relation $S_n^{[k-1]} = S_{n+1}^{[k]} - k S_n^{[k]}$ (3)

Proof: Using the expression (2), we obtain the following (see [8])

$$\begin{aligned} S_n^{[k-1]} &= \frac{1}{2\pi i} \oint_C \frac{z^{n-1}}{(z-1)\cdots(z-(k-1))} dz = \frac{1}{2\pi i} \oint_C \frac{z^n dz}{(z-1)\cdots(z-k)} - k \frac{1}{2\pi i} \oint_C \frac{z^{n-1} dz}{(z-1)\cdots(z-k)}, \\ &= S_{n+1}^{[k]} - k S_n^{[k]}, \quad q.e.d. \end{aligned}$$

There is a well-known generating function for the Stirling numbers of the second kind [8]. Upon using that generating function, (2) implies the property:

Manuscript received on 17 August 2024 | Revised Manuscript received on 29 August 2024 | Manuscript Accepted on 15 October 2024 | Manuscript published on 30 October 2024.


* Correspondence Author (s)

Dr. R. Sivaraman*, Department of Mathematics, Dwaraka Doss Goverdhan Doss Vaishnav College, Chennai 600 106, India. E-mail ID: rsivaraman1729@yahoo.co.in, ORCID ID: [0000-0001-5989-4422](https://orcid.org/0000-0001-5989-4422)

M. Muniru Iddrisu, Department of Mathematics, Faculty of Physical Sciences, University for Development Studies, Tamale, Ghana. E-mail ID: mmuniru@uds.edu.gh, ORCID ID: [0000-0001-7628-8168](https://orcid.org/0000-0001-7628-8168)

Prof. J. López-Bonilla, ESIME-Zacatenco, Instituto Politécnico Nacional, Edif. 4, 1er. Piso, Col. Lindavista CP 07738, CDMX, México. E-mail ID: jlopezb@ipn.mx, ORCID ID: [0000-0003-3147-7162](https://orcid.org/0000-0003-3147-7162)

© The Authors. Published by Lattice Science Publication (LSP). This is an open access article under the CC-BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>)



$$\sum_{m=0}^{\infty} S_m^{[n]} \frac{q^m}{m!} = \frac{1}{n!} (e^q - 1)^n = \frac{1}{2\pi i} \oint_C \frac{e^{zq} dz}{z(z-1)\cdots(z-n)}, \quad (4)$$

Further from (2), we can obtain the Following Compact Expression:

$$S_m^{[n]} = \frac{(-1)^n}{n!} (1 + \phi)^n, \quad \phi^k := (-1)^k k^m. \quad (5)$$

b).- Let us now consider the function $f(z) = \frac{x^z}{z!}$. Applying this function in (1), we obtain the Laguerre polynomials [10-15] as given below:

$$L_n(x) = \sum_{k=0}^n \binom{n}{k} (-1)^k \frac{x^k}{k!} = \frac{(-1)^n n!}{2\pi i} \oint_C \frac{\frac{x^z}{z!} dz}{z(z-1)\cdots(z-n)} = (1 + \beta)^n, \quad \beta^k := \frac{(-x)^k}{k!}, \quad (6)$$

From (6), we see that.

$$L_0(x) = 1, \quad L_1(x) = 1 - x, \quad L_2(x) = \frac{1}{2!} (2 - 4x + x^2), \quad L_3(x) = \frac{1}{3!} (6 - 18x + 9x^2 - x^3), \dots;$$

Also from (6), we can deduce the corresponding generating function:

$$\begin{aligned} \sum_{n=0}^{\infty} L_n(x) q^n &= \sum_{n=0}^{\infty} [(1 + \beta) q]^n = \frac{1}{1 - (1 + \beta) q} = \frac{1}{1 - q} \cdot \frac{1}{1 - \frac{q\beta}{1 - q}} = \frac{1}{1 - q} \left[1 + \frac{q\beta}{1 - q} + \left(\frac{q\beta}{1 - q} \right)^2 + \dots \right], \quad (7) \\ &= \frac{1}{1 - q} \left[1 + \frac{q}{1 - q} \left(-\frac{x}{1!} \right) + \left(\frac{q}{1 - q} \right)^2 \left(\frac{x^2}{2!} \right) + \left(\frac{q}{1 - q} \right)^3 \left(-\frac{x^3}{3!} \right) + \dots \right] = \frac{1}{1 - q} e^{-\frac{qx}{1 - q}}, \end{aligned}$$

c).- Melzak's identity [6, 8, 16-19]:

We now consider the function. $f(x + y) = x \binom{x+n}{n} \sum_{k=0}^n \binom{n}{k} (-1)^k \frac{f(y-k)}{x+k}$, $x, y \in \mathbb{C}$, $n \geq 0$, $x \neq 0, -1, -2, \dots, -n$, (8) or any algebraic polynomial $f(t)$ up to degree n ;

Then from (1) and (8) we obtain

$$f(x + y) = x \binom{x+n}{n} \frac{(-1)^n n!}{2\pi i} \oint_C \frac{\frac{f(y-z)}{x+z}}{z(z-1)\cdots(z-n)} dz, \quad x \neq 0, -1, \dots, -n, \quad (9)$$

for example, from (9) with $f(t) = 1$ we obtain

$$1 = x \binom{x+n}{n} \frac{(-1)^n n!}{2\pi i} \oint_C \frac{\frac{1}{x+z}}{z(z-1)\cdots(z-n)} dz \stackrel{(2)}{=} \binom{x+n}{n} (-1)^n n! \sum_{j=n}^{\infty} \frac{(-1)^j}{x^j} S_j^{[n]},$$

which simplifies to Melzak's identity given by

$$\sum_{k=n}^{\infty} \frac{(-1)^j}{x^j} S_j^{[n]} = \frac{(-1)^n}{(x+1)(x+2)\cdots(x+n)}. \quad (10)$$

III. CONCLUSION

Using the Nörlund-Rice integral formula as in (1), we have considered some specific functions in three possible cases. In particular, by thinking, $f(z) = z^m$ We proved a recurrence relation satisfied by Stirling's numbers of the second kind, as stated in Theorem 2.1. Using the function $f(z) = \frac{x^z}{z!}$ we have obtained nice expressions related to Laguerre polynomials and we have also used this formula to determine the generating function for such polynomials. Finally by considering $f(x + y) = x \binom{x+n}{n} \sum_{k=0}^n \binom{n}{k} (-1)^k \frac{f(y-k)}{x+k}$ we have obtained the Melzak's Identity as described in (10). These results and connections will provide new insights and rich applications

of the famous Norlund-Rice Integral formula. By considering additional functions, as we have done in this paper, researchers can aim to obtain more compelling results in the future.

DECLARATION STATEMENT

After aggregating input from all authors, I must verify the accuracy of the following information as the article's author.

- **Conflicts of Interest/ Competing Interests:** Based on my understanding, this article has no conflicts of interest.



- **Funding Support:** This article has not been sponsored or funded by any organization or agency. The independence of this research is a crucial factor in affirming its impartiality, as it was conducted without any external influence.
- **Ethical Approval and Consent to Participate:** The data provided in this article is exempt from the requirement for ethical approval or participant consent.
- **Data Access Statement and Material Availability:** The adequate resources of this article are publicly accessible.
- **Author's Contributions:** The authorship of this article is contributed equally to all participating individuals.

REFERENCES

1. N. E. Nörlund, *Vorlesungen über differenzenrechnung*, Chelsea (1954).
2. D. E. Knuth, *The Art of Computer Programming*, vol. 3, Addison-Wesley, New York (1973).
3. P. Flajolet, R. Sedgewick, *Mellin transforms and asymptotics: Finite differences and Rice's integrals*,
4. Theoret. Comput. Sci. **144**, No. 1-2 (1995) 101-124.
5. P. Kirschenhofer, *A note on alternating sums*, Electron. J. Combin. **3**, No. 2 (1996) Paper R7
6. P. Kirschenhofer, P. J. Larcombe, *On a class of recursive-based binomial coefficient identities involving*
7. *Harmonic numbers*, Util. Math. **73** (2007) 105-115.
8. K. N. Boyadzhiev, *Notes on the binomial transform*, World Scientific, Singapore (2018).
9. Ch. Benjes, *Properties of Stirling numbers of the second kind*, Master's Thesis, Naval Postgraduate School,
10. Monterey, Calif.-USA (1971).
11. J. Quaintance, H. W. Gould, *Combinatorial identities for Stirling numbers*, World Scientific, Singapore
12. (2016).
13. V. Barrera-Figueroa, J. López-Bonilla, R. López-Vázquez, *On Stirling numbers of the second kind*,
14. Prespacetime Journal **8**, No. 5 (2017) 572-574.
15. E. N. Laguerre, *Sur l'intégrale $\int_x^\infty x^{-1} e^{-x} dx$* , Bull. Soc. Math. France **7** (1879) 72-81.
16. F. Pidduck, *Laguerre polynomials in quantum mechanics*, J. London Math. Soc. **4**, No. 1 (1929)
17. 163-166.
18. C. Lanczos, *Linear differential operators*, SIAM, Philadelphia, USA (1996).
19. M. E. H. Ismail, *Classical and quantum orthogonal polynomials in one variable*, Cambridge University
20. Press (2005).
21. J. López-Bonilla, V. M. Salazar del Moral, R. López-Vázquez, *On some identities for the Laguerre*
22. *Polynomials*, Prespacetime J. **8**, No. 10 (2017) 1251-1254.
23. X. Pan, X. Guo, *Some New Identities Involving Laguerre Polynomials*, AIMS Math. **6**, No. 11 (2021)
24. 12713-12717.
25. 16.Z. A. Melzak, *Problem 4458*, Amer. Math. Monthly **58**, No. 9 (1951) 636.
26. U. Abel, H. W. Gould, J. Quaintance, *New proofs of Melzak's identity*, arXiv: 1603.07006v1 [math. CO]
27. Mar (2016).
28. J. López-Bonilla, J. Yaljá Montiel-Pérez, S. Vidal-Beltrán, *On the Melzak and Wilf identities*, The J. of
29. Middle East & North Africa Sci. **3**, No. 11 (2017) 1-3.
30. M. Gessel, *On Miki's identity for Bernoulli numbers*, J. Number Theory **110**, No. 1 (2005) 75-82.

AUTHOR'S PROFILE




Dr. R. Sivaraman, an Associate Professor at Dwaraka Doss Goverdhan Doss Vaishnav College, Chennai, has 25 years of teaching experience at the College level. He was conferred the National Award for Popularising Mathematics among the masses in 2016 by the Department of Science and Technology, Government of India. He was conferred with the Indian National Science Academy (INSA) Teaching award for the year 2018. He has also received the State Government's Best Science Book Award in 2011 and 2012. He has provided more than 400 lectures conveying the beauty and

applications of Mathematics. He has published over 200 research papers and completed his Postdoctoral Research Fellowship and doctorate in science. He has written 32 books to popularise mathematics among the general public. He was a member of the Textbook Writing Committee of the Tamil Nadu School Education Department, responsible for preparing the revised mathematics textbook for the eleventh class, and the chairperson for the tenth class. He has won more than 75 prestigious awards for his distinguished service to mathematics. He has been offering free courses to college students from impoverished backgrounds for many years. Propagating the beauty and applications of mathematics to everyone was his life mission.



Mohammed Muniru Iddrisu is a Professor of Mathematics at the University for Development Studies, Ghana. He holds a PhD in Mathematics and has supervised several undergraduate and postgraduate theses. His research interests include Mathematical Analysis, Coding Theory, Cryptography, and Mathematical Statistics, with an emphasis on Inequalities and Special functions. He is a member of many professional bodies. He has published several papers in reputed international journals. His passion is conducting research in mathematics and training several young students to pursue a career in mathematical research.



Professor José Luis López-Bonilla is a Teacher and Researcher at National Polytechnic Institute, Mexico city, and his Ph D is in Theoretical Physics: Electrodynamics [motion of classical charged particles], Special and General Relativity [Petrov classification, Lorentz matrix, Lanczos potential, Embedding of Riemannian spaces, Newman-Penrose and 2-spinors formalisms], Quantum Mechanics [matrix elements for several potentials] and Classical Mechanics [constrained Hamiltonian systems]. Besides, he has strong interest in Mathematical Methods Applied to Engineering [for example, importance of the SVD method in 5G technology] and also in specific topics in Number Theory such as: Partitions, Representations of integers as sums of squares, Combinatorics, Bernoulli and Stirling numbers and Recurrence relations for arithmetical functions.

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the Lattice Science Publication (LSP)/ journal and/ or the editor(s). The Lattice Science Publication (LSP)/ journal and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

