



One Integer Point on Some Elliptic Curve to Have Rank 1

Safwan Aouira, Hasan Sankari, Ahmad Abdo*

Abstract: Walsh proved a necessary condition for some curves of the form $y^2 = x^3 - px$, with p is a prime, to have Mordell - Weil rank 1. In this paper, we extend Walsh's result to show that some curves have Mordell-Weil rank 1.

Keywords. Elliptic Curves, Integer Points, Rank of Elliptic Curve, Mordell-Weil Rank, and Diophantine Equation.

I. INTRODUCTION

Many authors have studied the relationship between an elliptic curve's rank and the size of its set of Integer points. Silverman [3] showed that the number of integer solutions to a cubic Thue equation $f(x, y) = m$ is bounded in terms of the rank of the Mordell - Weil group of the corresponding cubic curve $f(x, y) = mz^3$.

For a curve of the form

$$E: y^2 = x^3 - px$$

With p prime, it is known that the rank of the Mordell-Weil group is at most 2, which can be proved using the 2-descent methods in Silverman [3]. H. Daghigh & S. Didari [1] studied the rank of $E: y^2 = x^3 = pqx$. In particular, Walsh [2] gave a general condition for the rank by considering the number of integer points on E . Recently, Mina and Bacani [6] determined necessary conditions for the rank of the curve $y^2 = x^3 - 3pqx$ to be zero.

II. PRELIMINARIES

In this section, we introduce the objects we need later and collect some important facts about them. Let E be an elliptic curve defined by

$$E: y^2 = x^3 - px$$

With n integer. And as in [6], we define

$$\tilde{E}: y^2 = x^3 + 4px.$$

To compute the rank (E) as in [6], we study the solvability of a Diophantine equation and rely on the methods of Sankari and Abdo [4], [5].

Let p is a prime and $n = 1, 2, 3$. Let

$$E_n/\mathbb{Q}: y^2 = x^3 - p^n x$$

Proposition 2.1. (See [3] or [7]). Let $E_n: y^2 = x^3 - p^n x$ be an elliptic curve, $\text{rank}_{E_n}(\mathbb{Q}) = 1$ if and only if exactly one of the equations

$$-S^4 + p^n T^4 = U^2 \quad (1)$$

$$2S^4 + 2p^n T^4 = U^2 \quad (2)$$

Has a primitive solution, such that $\gcd(S, T) = \gcd(S, U) = \gcd(T, U) = 1$.

III. NEW RESULTS

Our principal result is the following. In the following Theorems, we consider the integer points on some curves to have Mordell-Weil rank 1. Theorem 3. 1. Let $n = 1, u$ be odd, v be even, $r > 1$ be an integer and p is a prime number.

- If $(-1, 2u) \in E_1(\mathbb{Q})$, then $p = 4u^2 + 1$ and $\text{rank}(E_1) = 1$.
- If $(v^2, v(v^2 - 1)) \in E_1(\mathbb{Q})$, then $p = 2v^2 - 1$ and $\text{rank}(E_1) = 1$.
- If $(pt^2, \frac{pr(r^2+3)t}{2}) \in E_1(\mathbb{Q})$ such that $\text{norm}(r + \sqrt{2}t) = -1$ and $p \equiv 5 \pmod{8}$, then $p = r^2 + 4$ and $\text{rank}(E_1(\mathbb{Q})) = 1$.

Proof. i. We have $(-1, 2u) \in E_1(\mathbb{Q})$ we easily find that $p = 4u^2 + 1$ and $p(1)^4 - (1)^4 = (2u)^2$, therefore the equation (1) in Proposition 2.1 has a primitive solution. And it is easy to see that the equation (2) in Proposition Therefore, the equation (2) in Proposition 2.1 does not have a primitive solution (by mod 4). It follows from Proposition 2.1 that $\text{rank}(E_1) = 1$.

ii. we have $(v^2, v(v^2 - 1)) \in E_1(\mathbb{Q})$ we easily find that $p = 2v^2 - 1$ and $2p(1)^4 + 2(1)^4 = (2m)^2$, therefore the equation (2) in Proposition 2.1 does not have a primitive solution. It is easy to see that equation (1) in Proposition 2.1 has a primitive solution (by mod 4). It follows from Proposition 2.1 that $\text{rank}(E_1) = 1$.

iii. we have $(pt^2, \frac{pr(r^2+3)t}{2}) \in E_1(\mathbb{Q})$ such that $\text{norm}(r + \sqrt{2}t) = -1$, we easily find that $p = r^2 + 4$ and $p \equiv 5 \pmod{8}$, therefor the equation (1) in Proposition 2.1 has a primitive solution (this solution is $(1, 1, 2m)$). And it is easy to see that the equation (2) in Proposition 2.1 does not have a primitive solution (mod 8). It follows

Manuscript received on 11 April 2026 | First Revised Manuscript received on 19 May 2026 | Second Revised Manuscript received on 03 September 2026 | Manuscript Accepted on 15 October 2026 | Manuscript published on 30 October 2026.


*Correspondence Author(s)

Dr. Safwan Aouira, Professor, Department of Mathematics, Faculty of Science, Aleppo University, Aleppo, Syria. safwanaouira@gmail.com.

Dr. Hasan Sankari, Professor, Department of Mathematics, Faculty of Science, Lattakia University, Lattakia, Syria. Email ID: hasansankari@gmail.com.

Dr. Ahmad Abdo*, Department of Mathematics, Faculty of Science, Aleppo University, Aleppo, Syria. Email ID: abdoahmad926@gmail.com, ahmad.abdo@alepuniu.edu.sy, ORCID ID: 0000-0002-1333-3619.

© The Authors. Published by Lattice Science Publication (LSP). This is an open-access article under the CC-BY-NC-ND license. <http://creativecommons.org/licenses/by-nc-nd/4.0/>



from Proposition 2.1 that $\text{rank}(E_1) = 1$.

Theorem 3. 2. Let $n = 2, v$ be even and p is a prime number.

If $(v^2, v(v^2 - 1)) \in E_1(\mathbb{Q})$ and $p \equiv -1 \pmod{8}$, then $p^2 = 2v^2 - 1$ and $\text{rank}(E_1) = 1$.

proof. we have $(v^2, v(v^2 - 1)) \in E_1(\mathbb{Q})$ We easily find that $p = 2v^2 - 1$ and $2p^2(1)^4 + 2(1)^4 = (2m)^2$ Therefore, the equation (2) in Proposition 2.1 has a primitive solution. It is easy to see that the equation (1) in Proposition 2.1 does not have a primitive solution (by mod 8). It follows from Proposition 2.1 that $\text{rank}(E_1) = 1$.

Theorem 3. 3. Let $n = 3, u, v$ are integers and p is a prime number.

If $(-v^2, uv) \in E_1(\mathbb{Q})$ and $p \equiv 5 \pmod{8}$, then $p^3 = u^2 + v^4$ and $\text{rank}(E_1) = 1$.

proof. we have $(-v^2, uv) \in E_1(\mathbb{Q})$ We easily find that $p^3 = u^2 + v^4$ and $p^3(1)^4 - (b)^4 = (a)^2$, therefore the equation (1) in Proposition 2.1 has a primitive solution. It is easy to see that equation (2) in Proposition 2.1 does not have a primitive solution (by mod 8). It follows from Proposition 2.1 that $\text{rank}(E_1) = 1$.

IV. CONCLUSION

By extending Walsh's theorem to curves with Mordell-Weil rank 1, this study fills a critical gap in the literature. It provides a more comprehensive framework for understanding the arithmetic properties of these algebraic structures.

ACKNOWLEDGMENTS

We thank everyone who provided helpful advice and suggestions.

DECLARATION STATEMENT

Some of the cited references are older and are noted explicitly as [1], [3] and [7]. However, these works remain significant for the current study, as they are pioneering in their fields.

After aggregating input from all authors, I must verify the accuracy of the following information as the article's author.

- **Conflicts of Interest/ Competing Interests:** Based on my understanding, this article has no conflicts of interest.
- **Funding Support:** This article has not been funded by any organizations or agencies. This independence ensures that the research is conducted objectively and free from external influence.
- **Ethical Approval and Consent to Participate:** The data provided in this article is exempt from the requirement for ethical approval or participant consent.
- **Data Access Statement and Material Availability:** The adequate resources of this article are publicly accessible.
- **Author's Contributions:** Conceptualisation: Ahmad Abdo, Hasan Sankari and Safwan Aouira; methodology: Ahmad Abdo; investigation and writing- review and editing: Ahmad Abdo; project administration, validation and visualisation: Safwan Aouira and Hasan Sankari;

resources, formal analysis, and writing - original draft: Ahmed Abdo.

REFERENCES

1. H. Daghigh & S. Didari. On the elliptic curves of the curves of the form $y^2 = x^3 - pqx$. Iranian Journal of Mathematical Sciences and Informatics, 10(2)(2015), 77-86. <https://arxiv.org/pdf/2401.00215>
2. P. G. Walsh, Maximal ranks and integer points on a family of elliptic curves, to appear in Glasnik Mat. <https://projecteuclid.org/journals/tokyo-journal-of-mathematics/volume-34/issue-2/Integer-Points-and-Independent-Points-on-the-Elliptic-Curve-y2x3/10.3836/tjm/1327931392.pdf>
3. H. J. Silverman & T. J. Tate, Rational Points on Elliptic Curves. Springer, Second Edition(2015). <https://link.springer.com/book/10.1007/978-3-319-18588-0>
4. H. Sankari and A. Abdo, On Polynomial Solution of Pell's Equation, Journal of Mathematics, Volume 2021. <https://ideas.repec.org/a/hin/jimath/5379284.html>
5. H. Sankari and A. Abdo, The polynomial solutions of Quadratic Diophantine Equations $X^2 - p(t)Y^2 + 2K(t)X + 2p(t)L(t)Y = 0$ Journal of Mathematics, Volume 2021. https://www.academia.edu/165236619/On_Polynomial_Solutions_of_Pell's_Equation
6. R.J.S.Mina & B.J.Bacania. Elliptic curves of type $y^2 = x^3 - 3pqx$ having ranks zero and one, 17(1)(2023), 67-76. <https://mjms.upm.edu.my/fullpaper/2023-March-17-1/J.%20B.%20Bacani-67-76.pdf>
7. S.W.Kim .Ranks of elliptic curves $y^2 = x^3 \pm 4px$. International Journal of algebra, 9(5)(2015), 205-211. <https://www.m-hikari.com/ija/ija-2015/ija-5-8-2015/p/kimIJA5-8-2015.pdf>

AUTHOR'S PROFILE



Dr. Safwan Aouira is a Syrian scholar who obtained his PhD from Humboldt University in Berlin in 1990. He was appointed Professor at the University of Aleppo in the Syrian Arab Republic in 2002. He has published several scientific research papers in Scopus-indexed journals, including the Basic and Applied Sciences - Scientific Journal of King Faisal University. He has also authored several scientific books at Syrian universities. His main research interests include algebra and algebraic geometry.



Dr. Hassan Sankari is Syrian and obtained his PhD from the University of Caen Normandy in Caen, France, in 1992. He was appointed Professor at the University of Latakia in the Syrian Arab Republic in 2020. He has published several international scientific research papers in Scopus-indexed journals (Journal of Mathematics, Arab Journal of Mathematical Sciences, International Journal of Mathematics and Mathematical Sciences). His main research interests include algebra, number theory, and algebraic numbers.



Dr. Ahmad Abdo is Syrian and began his PhD thesis in Mathematics (Algebra) at the University of Aleppo in 2023. His main research interests include Algebra, Number Theory, and Algebraic Geometry. Mr Abdo has participated in several local conferences in Syria. He has published two papers in the Journal of Mathematics, an international Scopus-indexed journal.

Disclaimer/Publisher's Note: The statements, opinions, and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the Publisher /Journal and/or the editor(s). The Publisher /Journal and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions, or products referred to in the content.

