



A Family of Cotangent-Polylogarithm Integrals, Harmonic Sums, and Level-Four Cyclotomy

K. Srinivasa Raghava, R. Sivaraman*



Abstract: For integers $v \geq 2$ and parameters $0 \leq \lambda \leq 1$, we study the family $\mathcal{J}_v(\lambda) := \int_0^{\pi/2} \cot x \operatorname{Li}_v\left(\frac{\lambda}{1+\cot x}\right) dx$. We prove absolute convergence, reduce the integral to a Dirichlet series with positive coefficients, and evaluate it in terms of a depth-two multiple polylogarithm, with $\omega = (1+i)/2$: $\mathcal{J}_v(\lambda) = \frac{\pi}{2} \operatorname{Im} \operatorname{Li}_v(\lambda\omega) - \operatorname{Re} \operatorname{Li}_{v,1}(\lambda\omega, \omega^{-1})$. We then derive an exact second-order recurrence for the Dirichlet coefficients, an Euler differential descent in the deformation parameter, a functional-difference equation for the weight generating function, and an exponentially convergent expansion in generalized harmonic numbers and zeta values. As consequences, the sequence $\mathcal{J}_v := \mathcal{J}_v(1)$ is strictly decreasing, $\mathcal{J}_v = \frac{\pi}{4} + \left(\frac{\pi}{4} - \frac{1}{2}\right) 2^{-v} + O(3^{-v})$, $v \rightarrow \infty$, and no nontrivial linear recurrence with constant coefficients in v can hold. At $\lambda = 1$, the family gives a concrete one-dimensional model for level-four cyclotomic periods built from harmonic sums.

Keywords: Polylogarithm, Multiple Polylogarithm, Cyclotomic Period, Harmonic Number, Euler Sum, Dirichlet Series.

I. INTRODUCTION AND STATEMENTS

The classical polylogarithm, its multiple variants, and their special values connect analysis, number theory, and mathematical physics. Panzer [1] develops identities and evaluations for multiple polylogarithms, while Xu and Wang [2] study the associated harmonic Euler sums. Au [9, 10] gives systematic methods for one-dimensional polylogarithmic integrals and iterated integrals at algebraic arguments. Related parametric integral evaluations appear in Sofo, Pain, and Scharaschkin [11]; Shalev [12] studies log-sine integrals and Apéry-like sums through polylogarithmic generating functions. These works provide context for the present family; the specific evaluation and coefficient identities below are proved directly.

For integers $v \geq 2$ and $0 \leq \lambda \leq 1$, define

$$\mathcal{J}_v(\lambda) := \int_0^{\pi/2} \cot x \operatorname{Li}_v\left(\frac{\lambda}{1+\cot x}\right) dx, \quad \mathcal{J}_v := \mathcal{J}_v(1)$$

$$0 < \frac{\lambda}{1+\cot x} < 1 \quad (0 < x < \pi/2, 0 < \lambda \leq 1),$$

Manuscript received on 10 September 2026 | First Revised Manuscript received on 18 September 2026 | Second Revised Manuscript received on 02 October 2026 | Manuscript Accepted on 15 October 2026 | Manuscript published on 30 October 2026.

*Correspondence Author(s)

K. Srinivasa Raghava, Research Associate, Department of Mathematics, Pie Mathematics Association, Chennai (Tamil Nadu), India. Email ID: srinivasaraghava@gmail.com, ORCID ID: [0000-0003-0021-6322](https://orcid.org/0000-0003-0021-6322).

Dr. R. Sivaraman*, Associate Professor, Department of Mathematics, Dwaraka Doss Goverdhan Doss Vaishnav College, Arumbakkam, Chennai (Tamil Nadu), India. Email ID: piesiva3435@gmail.com, rsivaraman1729@yahoo.co.in, ORCID ID: [0000-0001-5989-4422](https://orcid.org/0000-0001-5989-4422).

© The Authors. Published by Lattice Science Publication (LSP). This is an open-access article under the CC-BY-NC-ND license. <http://creativecommons.org/licenses/by-nc-nd/4.0/>

We use the principal branch of Li_v on $\mathbb{C} \setminus [1, \infty)$, with the standard power-series normalization [13, §25.12(ii)]. Since no branch cut is crossed in (1). We also set

$$\omega := \frac{1+i}{2}, \bar{\omega} = \frac{1-i}{2}, \omega^{-1} = 1-i, |\omega| = 2^{-1/2}. \quad (2)$$

For complex α, β such that the series converges absolutely, our depth-two convention is

$$\operatorname{Li}_{v,1}(\alpha, \beta) := \sum_{\mu \geq \kappa \geq 1} \frac{\alpha^\mu \beta^\kappa}{\mu^\nu \kappa}. \quad (3)$$

This is the nested-series convention for multiple polylogarithms in [10, §2.2]; see also [1, §1, Eq. (4)] for an alternative convention using increasing summation indices.

The main evaluation is as follows.

Theorem 1.

For every integer $v \geq 2$ and every $0 \leq \lambda \leq 1$, the integral $\mathcal{J}_v(\lambda)$ is absolutely convergent. Moreover,

$$\mathcal{J}_v(\lambda) = \sum_{\mu=1}^{\infty} \frac{\alpha_\mu \lambda^\mu}{\mu^v}, \quad (4)$$

where

$$\alpha_\mu := \int_0^1 \frac{\eta^{\mu-1}(1-\eta)}{1-2\eta+2\eta^2} d\eta > 0, \quad \mu \geq 1, \quad (5)$$

and

$$\mathcal{J}_v(\lambda) = \frac{\pi}{2} \operatorname{Im} \operatorname{Li}_v(\lambda\omega) - \operatorname{Re} \operatorname{Li}_{v,1}(\lambda\omega, \omega^{-1}). \quad (6)$$

In particular,

$$\mathcal{J}_v = \frac{\pi}{2} \operatorname{Im} \operatorname{Li}_v(\omega) - \operatorname{Re} \operatorname{Li}_{v,1}(\omega, \omega^{-1}). \quad (7)$$

The coefficients α_μ have a simple exact recurrence.

Theorem 2.

For every integer

$$\mu \geq 1, \alpha_\mu - 2\alpha_{\mu+1} + 2\alpha_{\mu+2} = \frac{1}{\mu(\mu+1)}. \quad (8)$$

also

$$\alpha_1 = \frac{\pi}{4}, \quad \alpha_2 = \frac{\pi}{4} - \frac{1}{2}, \quad (9)$$

and


$$0 < \alpha_\mu \leq \frac{2}{\mu(\mu+1)} \quad (\mu \geq 1). \quad (10)$$

For the differential descent, extend (1) to $v = 1$ using $\operatorname{Li}_1(u) = -\log(1-u)$ and to $v = 0$ using $\operatorname{Li}_0(u) = u/(1-u)$. The corresponding integrals and series (1) converge for $0 \leq \lambda \leq 1$; at $(v, \lambda) = (0, 1)$ the endpoint is understood by its finite limit. These extensions also follow from (10). Term-wise differentiation of the classical polylogarithm series [13, Eq. (25.12.10)] gives the following descent.

Theorem 3.

Let $\vartheta := \lambda \frac{d}{d\lambda}$.

For every integer $v \geq 1$



and every $0 \leq \lambda < 1$,

$$\vartheta \mathcal{I}_v(\lambda) = \mathcal{I}_{v-1}(\lambda), \quad (11)$$

where

$$\mathcal{I}_0(\lambda) = \frac{\lambda(\pi + 2(1-\lambda)\log(1-\lambda))}{2(\lambda^2 - 2\lambda + 2)}. \quad (12)$$

Consequently,

$$\vartheta^v \mathcal{I}_v(\lambda) = \mathcal{I}_0(\lambda), \quad (13)$$

and, for $0 < \lambda < 1$,

$$\mathcal{I}_v(\lambda) = \frac{1}{(v-1)!} \int_0^\lambda \frac{\mathcal{I}_0(\tau)}{\tau} \log^{v-1}\left(\frac{\lambda}{\tau}\right) d\tau. \quad (14)$$

At $\lambda = 0$, the right-hand side is defined by continuity and equals zero.

The weight generating function satisfies a functional-difference equation.

Proposition 4.

$$\text{Let } \Phi(\sigma) := \sum_{v=1}^{\infty} \mathcal{I}_v \sigma^{v-1} \quad (|\sigma| < 1). \quad (15)$$

Then

$$\Phi(\sigma) = \sum_{\mu=1}^{\infty} \frac{\alpha_\mu}{\mu - \sigma}, \quad (16)$$

which extends meromorphically to $\mathbb{C}$, with simple poles at $1, 2, 3, \dots$. Moreover, as an identity of meromorphic functions,

$$\begin{aligned} \Phi(\sigma) - 2\Phi(\sigma+1) + 2\Phi(\sigma+2) \\ = \frac{1 - \gamma - \psi(1-\sigma)}{\sigma(\sigma+1)} \\ - \frac{\pi}{2(\sigma+1)}, \end{aligned} \quad (17)$$

where ψ is the digamma function and γ is Euler's constant.

The connection between nested polylogarithmic sums and harmonic Euler sums is classical [1, 2]. The depth-two term in (7) admits the following explicit harmonic expansion, derived in Section III.

Theorem 5.

For every integer $v \geq 2$,

$$\begin{aligned} \mathcal{I}_v = \frac{\pi}{2} \text{ImLi}_v(\omega) + \sum_{j=2}^v \zeta(j) \text{ReLi}_{v+1-j}(\omega) \\ - \sum_{j=1}^v \sum_{\rho=1}^{\infty} \frac{H_\rho^{(j)} \text{Re}(\omega^\rho)}{\rho^{v+1-j}}, \end{aligned} \quad (18)$$

where

$$H_\rho^{(j)} := \sum_{m=1}^{\rho} \frac{1}{m^j}, \quad H_\rho^{(1)} = H_\rho.$$

since

$$\text{Re}(\omega^\rho) = 2^{-\rho/2} \cos\left(\frac{\pi\rho}{4}\right), \quad (19)$$

Formula (18) is exponentially convergent.

A final consequence concerns monotonicity and asymptotics in the weight.

Corollary 6.

The Dirichlet series

$$\mathcal{D}(s) := \sum_{\mu=1}^{\infty} \frac{\alpha_\mu}{\mu^s} \quad (20)$$

converges absolutely for $\text{Res} > -1$, is holomorphic there, and satisfies

$$(-1)^r \frac{d^r}{ds^r} \mathcal{D}(s) > 0 \quad (s > -1, r = 0, 1, 2, \dots).$$

In particular, the sequence $\mathcal{I}_v = \mathcal{D}(v)$ is strictly decreasing, $\mathcal{I}_v \downarrow \pi/4$, and

$$\mathcal{I}_v = \frac{\pi}{4} + \left(\frac{\pi}{4} - \frac{1}{2}\right) 2^{-v} + O(3^{-v}). \quad (21)$$

Moreover, no nontrivial linear recurrence with constant coefficients in v can hold for the sequence $(\mathcal{I}_v)_{v \geq 2}$.

Remark 7.

The cyclotomic interpretation applies to the specialization $\lambda = 1$. In the unit-interval variable η , the relevant singular points are $0, 1, \omega, \bar{\omega}, \infty$, since

$$1 - 2\eta + 2\eta^2 = (1 - \eta/\omega)(1 - \eta/\bar{\omega}).$$

The Möbius transformation $\tau = \eta/(1 - \eta)$ sends these points to $0, \infty, i, -i, -1$, respectively, a subset of the level-four alphabet $\{0, 1, -1, i, -i, \infty\}$. Thus, the iterated integrals for $\mathcal{I}_v(1)$ have a level-four cyclotomic description. The invariance under Möbius changes of variables and the identification with cyclotomic multiple zeta values are made precise in [10, Proposition 3.2(4), Theorem 3.5, and Corollary 3.6].

For the underlying cyclotomic theory, see Goncharov [3, §§1.1 and 2.8]; for double-shuffle and other standard relations at roots of unity, see Yaddaden [4] and Bachmann and Yaddaden [6]. Brown provides the broader motivic framework [5, §10.6.2], while Ablinger and Schneider [7] study cyclotomic harmonic sums, and Ablinger, Blümlein, Round, and Schneider [8, §2] review cyclotomic polylogarithms. These references concern the structural setting, not the particular evaluation (6).

For $0 < \lambda < 1$, $\text{Li}_v(\lambda\eta)$ introduces the additional point $\eta = 1/\lambda$, which maps to $\tau = 1/(\lambda - 1)$. Consequently, the same argument does not establish a level-four period statement for arbitrary λ ; the general family remains polylogarithmic, with a parameter-dependent alphabet.

II. PROOF OF THE EVALUATION

We begin with convergence and the reduction to the unit interval.

Lemma 8.

For $v \geq 2$ and $0 \leq \lambda \leq 1$,

$$\mathcal{I}_v(\lambda) = \int_0^1 \text{Li}_v(\lambda\eta) \frac{1 - \eta}{\eta(1 - 2\eta + 2\eta^2)} d\eta. \quad (22)$$

The integral is absolutely convergent.

Proof. As $x \rightarrow 0^+$,

$$\cot x = \frac{1}{x} + O(x), \quad \frac{\lambda}{1 + \cot x} = \lambda x + O(x^2),$$

$$\text{Li}_v(u) = u + O(u^2),$$

So the integrand in (1) equals $\lambda + O(x)$. As $x \rightarrow \pi/2^-$, if $u := \pi/2 - x$, then

$$\cot x = u + O(u^3), \quad \frac{\lambda}{1 + \cot x} = \lambda + O(u),$$

and $\text{Li}_v(\lambda + O(u))$ remains bounded for $0 \leq \lambda \leq 1$, so the integrand is $O(u)$. Hence the integral is absolutely convergent. Now set $\tau = \tan x$ and then $\eta = \tau/(1 + \tau)$.





Since

$$\frac{\lambda}{1 + \cot x} = \frac{\lambda \tau}{1 + \tau} = \lambda \eta, \quad dx = \frac{d\tau}{1 + \tau^2} = \frac{d\eta}{1 - 2\eta + 2\eta^2},$$

and $\cot x dx = \frac{1-\eta}{\eta(1-2\eta+2\eta^2)} d\eta$,
formula (22) follows.

Lemma 9.

For $v \geq 2$ and $0 \leq \lambda \leq 1$,

$$\mathcal{J}_v(\lambda) = \sum_{\mu=1}^{\infty} \frac{\alpha_{\mu} \lambda^{\mu}}{\mu^v},$$

$$\alpha_{\mu} = \int_0^1 \frac{\eta^{\mu-1}(1-\eta)}{1-2\eta+2\eta^2} d\eta.$$

In particular, each α_{μ} is positive.

Proof. For $0 \leq \eta \leq 1$ and $0 \leq \lambda \leq 1$, the defining polylogarithm series [13, Eq. (25.12.10)]

$$\text{Li}_v(\lambda \eta) = \sum_{\mu=1}^{\infty} \frac{(\lambda \eta)^{\mu}}{\mu^v}$$

has nonnegative terms. By Tonelli's theorem and Lemma 8,

$$\mathcal{J}_v(\lambda) = \sum_{\mu=1}^{\infty} \frac{\lambda^{\mu}}{\mu^v} \int_0^1 \frac{\eta^{\mu-1}(1-\eta)}{1-2\eta+2\eta^2} d\eta,$$

which is the claim.

Lemma 10.

For every integer $\mu \geq 1$,

$$\alpha_{\mu} = \frac{\pi}{2} \text{Im}(\omega^{\mu}) - \text{Re} \sum_{\kappa=1}^{\mu-1} \frac{\omega^{\mu-\kappa}}{\kappa}. \quad (23)$$

Proof. Because

$$1 - 2\eta + 2\eta^2 = (1 - \eta/\omega)(1 - \eta/\bar{\omega}),$$

a direct partial fraction decomposition gives

$$\frac{1-\eta}{\eta(1-2\eta+2\eta^2)} = \frac{1}{\eta} + \frac{\bar{\omega}}{1-\eta/\omega} + \frac{\omega}{1-\eta/\bar{\omega}}. \quad (24)$$

Hence

$$\alpha_{\mu} = \frac{1}{\mu} + \bar{\omega} \mathcal{J}_{\mu}(\omega^{-1}) + \omega \mathcal{J}_{\mu}(\bar{\omega}^{-1}) \quad (24a)$$

where

$$\mathcal{J}_{\mu}(\alpha) := \int_0^1 \frac{\eta^{\mu}}{1-\alpha\eta} d\eta.$$

If $\alpha \notin [1, \infty)$ and $\alpha \neq 0$, the segment from 0 to α does not cross the principal logarithmic cut, and the substitution $u = \alpha\eta$ yields

$$\mathcal{J}_{\mu}(\alpha) = \alpha^{-\mu-1} \int_0^{\alpha} \frac{u^{\mu}}{1-u} du.$$

Using

$$\frac{u^{\mu}}{1-u} = \frac{1}{1-u} - \sum_{\kappa=0}^{\mu-1} u^{\kappa},$$

we obtain

$$\mathcal{J}_{\mu}(\alpha) = -\alpha^{-\mu-1} \left(\log(1-\alpha) + \sum_{\kappa=1}^{\mu} \frac{\alpha^{\kappa}}{\kappa} \right). \quad (25)$$

Now

$$1 - \omega^{-1} = i, \quad 1 - \bar{\omega}^{-1} = -i, \quad \log(i) = \frac{i\pi}{2}, \quad \log(-i) = -\frac{i\pi}{2},$$

with the principal logarithm. Substituting (25) into (24a), using $\bar{\omega}\omega^{\mu+1} = \omega^{\mu}/2$ and $\omega\bar{\omega}^{\mu+1} = \bar{\omega}^{\mu}/2$, and then cancelling the term $1/\mu$, we get

$$\alpha_{\mu} = \frac{\pi}{2} \text{Im}(\omega^{\mu}) - \text{Re} \sum_{\kappa=1}^{\mu-1} \frac{\omega^{\mu-\kappa}}{\kappa},$$

as required.

Proof of Theorem 1. Lemma 8 gives absolute convergence, and Lemma 9 gives (4). Multiply (23) by λ^{μ}/μ^v and sum over $\mu \geq 1$. The first term gives

$$\sum_{\mu=1}^{\infty} \frac{\lambda^{\mu}}{\mu^v} \frac{\pi}{2} \text{Im}(\omega^{\mu}) = \frac{\pi}{2} \text{Im} \sum_{\mu=1}^{\infty} \frac{(\lambda\omega)^{\mu}}{\mu^v} = \frac{\pi}{2} \text{Im} \text{Li}_v(\lambda\omega).$$

For the second term we write $\rho = \mu - \kappa \geq 1$ and estimate

$$\sum_{\mu > \kappa \geq 1} \frac{|\lambda\omega|^{\mu} |\omega|^{-\kappa}}{\mu^v \kappa} = \sum_{\rho=1}^{\infty} |\omega|^{\rho} \sum_{\kappa=1}^{\infty} \frac{\lambda^{\rho+\kappa}}{(\rho+\kappa)^v \kappa}.$$

The inner sum is

$$\ll \rho^{-v} \sum_{\kappa \leq \rho} \frac{1}{\kappa} + \sum_{\kappa > \rho} \frac{1}{\kappa^{v+1}} \ll \rho^{-v} \log(\rho+1),$$

So the outer sum converges because $|\omega| < 1$. Therefore, the double series converges absolutely, and we may rearrange:

$$\sum_{\mu=1}^{\infty} \frac{\lambda^{\mu}}{\mu^v} \text{Re} \sum_{\kappa=1}^{\mu-1} \frac{\omega^{\mu-\kappa}}{\kappa} = \text{Re} \sum_{\mu > \kappa \geq 1} \frac{(\lambda\omega)^{\mu} \omega^{-\kappa}}{\mu^v \kappa} = \text{Re} \text{Li}_{v,1}(\lambda\omega, \omega^{-1}).$$

This proves (6). The special case (7) is obtained by setting $\lambda = 1$.

III. STRUCTURAL CONSEQUENCES

Proof of Theorem 2. The bound follows from

$$1 - 2\eta + 2\eta^2 = 2 \left(\eta - \frac{1}{2} \right)^2 + \frac{1}{2} \geq \frac{1}{2}, \quad 0 \leq \eta \leq 1,$$

which gives

$$0 < \alpha_{\mu} \leq 2 \int_0^1 \eta^{\mu-1} (1-\eta) d\eta = \frac{2}{\mu(\mu+1)}.$$

also,

$$\alpha_{\mu} - 2\alpha_{\mu+1} + 2\alpha_{\mu+2} = \int_0^1 \eta^{\mu-1} (1-\eta) d\eta = \frac{1}{\mu(\mu+1)},$$

which is (8). Finally,

$$\alpha_1 = \int_0^1 \frac{1-\eta}{1-2\eta+2\eta^2} d\eta = \frac{\pi}{4},$$

$$\alpha_2 = \int_0^1 \frac{\eta(1-\eta)}{1-2\eta+2\eta^2} d\eta = \frac{\pi}{4} - \frac{1}{2}.$$

Proof of Theorem 3. For $0 \leq \lambda < 1$, the series in (4) converges locally uniformly in λ ; hence termwise differentiation is valid and gives

$$\vartheta \mathcal{J}_v(\lambda) = \sum_{\mu=1}^{\infty} \frac{\alpha_{\mu} \lambda^{\mu}}{\mu^{v-1}} = \mathcal{J}_{v-1}(\lambda).$$

To compute $\mathcal{J}_0(\lambda)$, start from Lemma 8 with $\text{Li}_0(u) = u/(1-u)$:

$$\mathcal{J}_0(\lambda) = \int_0^1 \frac{\lambda(1-\eta)}{(1-\lambda\eta)(1-2\eta+2\eta^2)} d\eta.$$

A direct decomposition yields



$$\frac{\lambda(1-\eta)}{(1-\lambda\eta)(1-2\eta+2\eta^2)} = \frac{\lambda^2(\lambda-1)}{\lambda^2-2\lambda+2} \frac{1}{1-\lambda\eta} + \frac{\lambda(\lambda-1)}{\lambda^2-2\lambda+2} \frac{1}{1-2\eta+2\eta^2} + \frac{\lambda}{\lambda^2-2\lambda+2} \frac{1}{1-2\eta+2\eta^2}$$

The middle term integrates to zero, while

$$\int_0^1 \frac{d\eta}{1-2\eta+2\eta^2} = \frac{\pi}{2}.$$

This gives (12). Formula (13) follows by iteration. Finally, (14) is the unique solution of (11) with $\mathcal{I}_v(0) = 0$, obtained by integrating the first-order equation repeatedly.

Proof of Proposition 4. From (10), we have $\alpha_\mu = O(\mu^{-2})$, so the series in (16) converges locally uniformly on compact subsets of $\mathbb{C} \setminus \{1, 2, 3, \dots\}$. For $|\sigma| < 1$,

$$\sum_{v=1}^{\infty} \frac{\sigma^{v-1}}{\mu^v} = \frac{1}{\mu - \sigma},$$

and therefore

$$\Phi(\sigma) = \sum_{\mu=1}^{\infty} \alpha_\mu \sum_{v=1}^{\infty} \frac{\sigma^{v-1}}{\mu^v} = \sum_{\mu=1}^{\infty} \frac{\alpha_\mu}{\mu - \sigma}.$$

Next,

$$\begin{aligned} \Phi(\sigma+1) &= -\frac{\alpha_1}{\sigma} + \sum_{\mu=1}^{\infty} \frac{\alpha_{\mu+1}}{\mu - \sigma}, \Phi(\sigma+2) \\ &= -\frac{\alpha_1}{\sigma+1} - \frac{\alpha_2}{\sigma} + \sum_{\mu=1}^{\infty} \frac{\alpha_{\mu+2}}{\mu - \sigma}. \end{aligned}$$

Using Theorem 2,

$$\begin{aligned} \Phi(\sigma) - 2\Phi(\sigma+1) + 2\Phi(\sigma+2) \\ = \sum_{\mu=1}^{\infty} \frac{1}{\mu(\mu+1)(\mu-\sigma)} + \frac{1}{\sigma} - \frac{\pi}{2(\sigma+1)}. \end{aligned}$$

Now

$$\begin{aligned} \frac{1}{\mu(\mu+1)(\mu-\sigma)} &= \frac{1}{\sigma(\sigma+1)} \left(\frac{1}{\mu-\sigma} - \frac{1}{\mu} \right) \\ &\quad - \frac{1}{\sigma+1} \left(\frac{1}{\mu} - \frac{1}{\mu+1} \right), \end{aligned}$$

so the classical digamma expansion [13, Eq. (5.7.6)]

$$-\psi(1-\sigma) - \gamma = \sum_{\mu=1}^{\infty} \left(\frac{1}{\mu-\sigma} - \frac{1}{\mu} \right)$$

yields

$$\sum_{\mu=1}^{\infty} \frac{1}{\mu(\mu+1)(\mu-\sigma)} = \frac{-\psi(1-\sigma) - \gamma}{\sigma(\sigma+1)} - \frac{1}{\sigma+1}.$$

Substitution proves (17).

Proof of Theorem 5. The reduction below uses the harmonic-sum language familiar from the study of Euler sums; see Xu and Wang [2] for that broader context. Start from (7) and write $\rho = \mu - \kappa$ in the depth-two series:

$$\text{Li}_{v,1}(\omega, \omega^{-1}) = \sum_{\rho=1}^{\infty} \omega^\rho \sum_{\kappa=1}^{\infty} \frac{1}{\kappa(\kappa+\rho)^v}.$$

For fixed $\rho \geq 1$ and $v \geq 2$, a direct partial fraction identity gives

$$\frac{1}{\kappa(\kappa+\rho)^v} = \frac{1}{\rho^v} \left(\frac{1}{\kappa} - \frac{1}{\kappa+\rho} \right) - \sum_{j=2}^v \frac{1}{\rho^{v+1-j}(\kappa+\rho)^j}. \quad (26)$$

Summing over $\kappa \geq 1$, we obtain

$$\sum_{\kappa=1}^{\infty} \frac{1}{\kappa(\kappa+\rho)^v} = \frac{H_\rho}{\rho^v} - \sum_{j=2}^v \frac{\zeta(j) - H_\rho^{(j)}}{\rho^{v+1-j}}.$$

Therefore

$$\text{Li}_{v,1}(\omega, \omega^{-1}) = \sum_{\rho=1}^{\infty} \omega^\rho \left(\frac{H_\rho}{\rho^v} - \sum_{j=2}^v \frac{\zeta(j) - H_\rho^{(j)}}{\rho^{v+1-j}} \right).$$

Take real parts and use

$$\sum_{\rho=1}^{\infty} \frac{\text{Re}(\omega^\rho)}{\rho^s} = \text{ReLi}_s(\omega), \quad \text{Re}(\omega^\rho) = 2^{-\rho/2} \cos\left(\frac{\pi\rho}{4}\right),$$

which is (19). Formula (18) follows.

Proof of Corollary 6. Because of (10), the series (20) converges absolutely for $\text{Re}s > -1$ and defines a holomorphic function there. For real $s > -1$ and integers $r \geq 0$, termwise differentiation gives

$$(-1)^r \frac{d^r}{ds^r} \mathcal{D}(s) = \sum_{\mu=1}^{\infty} \frac{\alpha_\mu (\log \mu)^r}{\mu^s} > 0,$$

so $\mathcal{D}$ is completely monotone on $(-1, \infty)$. In particular, $\mathcal{I}_{v+1} < \mathcal{I}_v$.

From Theorem 2,

$$\mathcal{I}_v = \alpha_1 + \alpha_2 2^{-v} + \sum_{\mu=3}^{\infty} \frac{\alpha_\mu}{\mu^v} = \frac{\pi}{4} + \left(\frac{\pi}{4} - \frac{1}{2} \right) 2^{-v} + R_v.$$

Using (10),

$$0 < R_v \leq 3^{-v} \sum_{\mu=3}^{\infty} \alpha_\mu \leq 2 \cdot 3^{-v} \sum_{\mu=3}^{\infty} \frac{1}{\mu(\mu+1)} = \frac{2}{3} 3^{-v},$$

which proves (21) and $\mathcal{I}_v \rightarrow \alpha_1 = \pi/4$.

Finally, assume that constants $c_0, \dots, c_r$ satisfy

$$\sum_{j=0}^r c_j \mathcal{I}_{v+j} = 0 \quad \text{for all large } v.$$

Using (4) with $\lambda = 1$, we obtain

$$\sum_{\mu=1}^{\infty} \alpha_\mu \mu^{-v} \left(\sum_{j=0}^r c_j \mu^{-j} \right) = 0 \quad \text{for all large } v.$$


Set $P(z) := \sum_{j=0}^r c_j z^j$. If some coefficient $a_\mu := \alpha_\mu P(1/\mu)$ is nonzero, let m be the least such index. Multiplication by m^v gives

$$0 = a_m + \sum_{\mu > m} a_\mu \left(\frac{m}{\mu} \right)^v.$$

Since P is bounded on $[0, 1]$ and $\sum_\mu \alpha_\mu < \infty$ by (10), the sequence (a_μ) is absolutely summable. Dominated convergence as $v \rightarrow \infty$ therefore gives $a_m = 0$, a contradiction. Thus, using $\alpha_\mu > 0$,

$$\sum_{j=0}^r c_j \mu^{-j} = 0 \quad \text{for every } \mu \geq 1.$$

A nonzero polynomial in $1/\mu$ cannot vanish at infinitely many points, so all c_j vanish.





Remark 11.

The involution $\tau \mapsto \tau^{-1}$ in the form

$$\mathcal{I}_v = \int_0^\infty \frac{\text{Li}_v\left(\frac{\tau}{1+\tau}\right)}{\tau(1+\tau^2)} d\tau$$

gives the reciprocal identity

$$\mathcal{I}_v = \int_0^\infty \frac{\tau \text{Li}_v\left(\frac{1}{1+\tau}\right)}{1+\tau^2} d\tau.$$

This symmetry is elementary, but it fits well with the relation $\omega/(1-\omega) = i$ behind the level-four factorization. The factors $1+\tau$ and $1+\tau^2$ are the cyclotomic polynomials $\Phi_2(\tau)$ and $\Phi_4(\tau)$, respectively; their rational kernels are part of the cyclotomic-polylogarithm alphabet in [8, §2, Eqs. (2.1)–(2.2)].

IV. CONCLUSION

We obtained exact polylogarithmic representations of the cotangent integral family. Its Dirichlet coefficients yield recurrence, differential, and generating-function identities. At $\lambda = 1$, an exponentially convergent harmonic expansion complements the weight asymptotics. The integral values decrease to $\pi/4$ and satisfy no nontrivial constant-coefficient linear recurrence in the weight. This specialization admits a level-four cyclotomic interpretation. For $0 < \lambda < 1$, the argument does not establish the same interpretation.

DECLARATION STATEMENT

After aggregating input from all authors, I must verify the accuracy of the following information as the article's author.

- **Conflicts of Interest/ Competing Interests:** Based on my understanding, this article has no conflicts of interest.
- **Funding Support:** This article has not been funded by any organisations or agencies. This independence ensures that the research is conducted objectively and free from external influence.
- **Ethical Approval and Consent to Participate:** The data provided in this article is exempt from the requirement for ethical approval or participant consent.
- **Data Access Statement and Material Availability:** The data and materials in this article are publicly accessible.
- **Authors' Contributions:** All authors contributed equally to this article.

REFERENCES

1. E. Panzer, The parity theorem for multiple polylogarithms, *J. Number Theory* 172 (2017), 93–113. DOI: [10.1016/j.jnt.2016.08.004](https://doi.org/10.1016/j.jnt.2016.08.004).
2. C. Xu and W. Wang, Explicit formulas of Euler sums via multiple zeta values, *J. Symbolic Comput.* 101 (2020), 109–127. DOI: [10.1016/j.jsc.2019.06.009](https://doi.org/10.1016/j.jsc.2019.06.009).
3. A. B. Goncharov, Motivic fundamental group of $\mathbb{G}_m - \mu_N$ and modular manifolds, *arXiv:1910.10321 [math.AG]*, 2019. <https://arxiv.org/abs/1910.10321>.
4. K. Yaddaden, The cyclotomic double shuffle torsor in terms of Betti and de Rham coproducts, *J. Pure Appl. Algebra* 228 (2024), no. 5, Paper No. 107550, 66 pp. DOI: [10.1016/j.jpaa.2023.107550](https://doi.org/10.1016/j.jpaa.2023.107550).
5. F. Brown, Notes on motivic periods, *Commun. Number Theory Phys.* 11 (2017), no. 3, 557–655. DOI: [10.4310/CNTP.2017.v11.n3.a2](https://doi.org/10.4310/CNTP.2017.v11.n3.a2).
6. H. Bachmann and K. Yaddaden, On a conjecture of Zhao related to standard relations among cyclotomic multiple zeta values, *J. Algebra* 688 (2026), 21–58. DOI: [10.1016/j.jalgebra.2025.10.003](https://doi.org/10.1016/j.jalgebra.2025.10.003).

7. J. Ablinger and C. Schneider, Algebraic independence of sequences generated by (cyclotomic) harmonic sums, *Ann. Comb.* 22 (2018), 213–244. DOI: [10.1007/s00026-018-0381-5](https://doi.org/10.1007/s00026-018-0381-5).
8. J. Ablinger, J. Blümlein, M. Round, and C. Schneider, Special functions, transcendentals and their numerics, *PoS(RADCOR2017)* (2018), 010. DOI: [10.22323/1.290.0010](https://doi.org/10.22323/1.290.0010).
9. K. C. Au, Evaluation of one-dimensional polylogarithmic integral, with applications to infinite series, *arXiv:2007.03957 [math.NT]*, 2020; version 4, 29 January 2024. <https://arxiv.org/abs/2007.03957v4>.
10. K. C. Au, Iterated integrals and multiple polylogarithm at algebraic arguments, *arXiv:2201.01676 [math.NT]*, 2022; version 4, 18 August 2026. <https://arxiv.org/abs/2201.01676v4>.
11. A. Sofo, J.-C. Pain, and V. Scharaschkin, A family of polylogarithmic integrals, *J. Appl. Anal.* 32 (2026), no. 1, 25–34; published online 13 January 2025. DOI: [10.1515/jaa-2024-0133](https://doi.org/10.1515/jaa-2024-0133). Preprint: <https://arxiv.org/abs/2412.02031>.
12. N. Shalev, Closed-form evaluations of log-sine integrals and Apéry-like sums in terms of polylogarithms, *Integral Transforms Spec. Funct.* (2026), published online 5 July 2026, 1–23. DOI: [10.1080/10652469.2026.2698652](https://doi.org/10.1080/10652469.2026.2698652). Preprint: <https://arxiv.org/abs/2508.05770>.
13. National Institute of Standards and Technology, NIST Digital Library of Mathematical Functions, version 1.2.8 (15 September 2026), §25.12 (polylogarithms), especially Eq. (25.12.10), and Eq. (5.7.6) (digamma series). <https://dlmf.nist.gov/>. Accessed 15 October 2026.

AUTHORS' PROFILE



K. Srinivasa Raghava is a mathematics researcher in number theory, cryptography, graph theory, and work inspired by Srinivasa Ramanujan. A mathematics and Vedic Math teacher and a chess player, he is a member of the MAA and AIRMC. He received the Prathibha Shiromani Award, the Math Genius Award, and the A.P.J. Abdul Kalam

Awards. He has presented papers and delivered invited talks at 72 national and 65 international conferences. He is passionate about creating new and exciting formulas in mathematics and extending mathematical concepts with philosophy, quantum mechanics, and computer science. His posts about mathematics on various social media platforms have influenced many young students and mathematics enthusiasts worldwide.



Dr R. Sivaraman, working as an Associate Professor at Dwaraka Doss Goverdhan Doss Vaishnav College, Chennai, has more than 28 years of teaching experience at the college level. In 2016, the Department of Science and Technology, Government of India, conferred on him the National Award for Popularising Mathematics among the masses. He received the Indian National Science Academy (INSA) Teaching Award in 2018. He has also received State Government Best Science Book Awards in 2011 and 2012. He has provided more than 500 lectures conveying the beauty and applications of Mathematics. He has published more than 300 research papers and has completed his Postdoctoral Research Fellowship and Doctor of Science Degree. For his extensive research spanning several branches of science, Stanford University and Elsevier Publications have included his name in the Top 2% Scientists list for 2025. He has written 53 books to popularise mathematics among the common man. He served on the Textbook Writing Committee of the Tamil Nadu School Education Department to prepare a revised mathematics textbook for eleventh class, and chaired the committee for tenth class. He has won more than 75 prestigious awards for his distinguished service to mathematics. He has taught free classes to college students from very poor backgrounds for many years. His life mission was to spread the beauty and applications of mathematics to everyone.

Disclaimer/Publisher's Note: The statements, opinions, and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the Publisher/Journal and/or the editor(s). The Publisher/Journal and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions, or products referred to in the content.

